arXiv:hep-ph/0702239v2 7 Apr 2007

Relativistic corrections to J/ψ exclusive and inclusive double charm production at B factories

Zhi-Guo He,[1,*] Ying Fan,[1,†] and Kuang-Ta Chao[1,2,‡]

[1]*Department of Physics, Peking University, Beijing 100871, China*

[2]*China Center of Advanced Science and Technology (World Laboratory), Beijing 100080, China*

Abstract

In order to clarify the puzzling problems in double charm production, relativistic corrections at order v^2 to the processes $e^+e^- \to J/\psi + \eta_c$ and $e^+e^- \to J/\psi + c\overline{c}$ at B factories are studied in non-relativistic quantum chromodynamics. The short-distance parts of production cross sections are calculated perturbatively, while the long-distance matrix elements are estimated from J/ψ and η_c decays up to errors of order v^4. Our results show that the relativistic correction to the exclusive process $e^+e^- \to J/\psi+\eta_c$ is significant, which, when combined together with the next-to-leading order α_s corrections, could resolve the large discrepancy between theory and experiment; whereas for the inclusive process $e^+e^- \to J/\psi+c\overline{c}$ the relativistic correction is tiny and negligible. The physical reason for the above difference between exclusive and inclusive processes largely lies in the fact that in the exclusive process the relative momentum between quarks in charmonium substantially reduces the virtuality of the gluon that converts into a charm quark pair, but this is not the case for the inclusive process, in which the charm quark fragmentation $c \to J/\psi + c$ is significant, and QCD radiative corrections can be more essential.

PACS numbers: 12.38.Cy, 13.66.Bc, 14.40.Gx

*Electronic address: hzgzlh@gmail.com

†Electronic address: fanying@pku.edu.cn

‡Electronic address: ktchao@th.phy.pku.edu.cn

I. INTRODUCTION

The exclusive and inclusive double charm production in e^+e^- annihilation at B factories has been one of the most puzzling problems in nonrelativistic QCD (NRQCD) [1] and heavy quarkonium physics [2] for years. The measured cross sections for exclusive double charmonium production $e^+e^- \to J/\psi + \eta_c$ and inclusive double charm production $e^+e^- \to J/\psi + c\bar{c}$ at $\sqrt{s} = 10.6$ GeV are much higher than the leading order (LO) predictions in NRQCD.

The cross section of $e^+e^- \to J/\psi + \eta_c$ at $\sqrt{s} = 10.6$ GeV observed by Belle[3, 4] is

$$\sigma[e^+e^- \to J/\psi + \eta_c] \times \mathcal{B}^{\eta_c}[\geq 2] = 25.6 \pm 2.8 \pm 3.4 \ fb, \tag{1a}$$

and by BaBar[5] is

$$\sigma[e^+e^- \to J/\psi + \eta_c] \times \mathcal{B}^{\eta_c}[\geq 2] = 17.6 \pm 2.8 \pm 2.1 \ fb, \tag{1b}$$

where $\mathcal{B}^{\eta_c}[\geq 2]$ is the branching fraction for η_c decay into at least two charged particles. Since $\mathcal{B}^{\eta_c}[\geq 2] < 1$, Eq.[1] is the lower bound of the cross section of the double charmonium production. It is about an order of magnitude larger than the theoretical predictions[6, 7, 8], at the leading order of strong coupling constant α_s and quark relative velocity v in the NRQCD factorization approach[1]. In[9] the next-to leading order (NLO) QCD result is given and it is found that the $\mathcal{O}(\alpha_s)$ corrections may greatly reduce the large discrepancy between theory and experiment. Meanwhile, the relativistic corrections are also considered by a number of authors. In [10], relativistic and bound state effects are discussed on the basis of relativistic quark models. In [11, 12, 13, 14, 15], relativistic effects are estimated in the framework of the light cone method, and very large enhancement effects on the cross sections can be found with certain light-cone distribution amplitudes. In [16], the authors calculate the relativistic correction based on NRQCD and light cone method with long-distance matrix elements determined from the Cornell potential model, and they point out, however, that after subtracting parts of the light-cone distribution functions that correspond to corrections of relative-order α_s in NRQCD, the enhancement effect due to relativistic corrections is not very large but still substantial. Probably, with both relative-order-α_s and relativistic corrections we may resolve the discrepancy between theory and experiment for $e^+e^- \to J/\psi + \eta_c$[16] .

The large discrepancies between theory and experiment exist not only in the exclusive

double charmonium production processes such as $e^+e^- \to J/\psi + \eta_c$ but also in the inclusive production process of $e^+e^- \to J/\psi + c\bar{c}$. The experimental result measured by Belle[3] is

$$\sigma[e^+e^- \to J/\psi + c\bar{c} + X] = 0.87^{+0.21}_{-0.19} \pm 0.17\text{pb}, \tag{2}$$

which is about five times larger than the theoretical calculations[17, 18, 19, 20, 21] based on NRQCD at leading order of α_s and v^2. In [22] the two-photon contribution and in [21] the color-octet contribution to $J/\psi + c\bar{c}$ production are further considered, but they are too small to resolve the large discrepancy. Other attempts to solve the problem can be found in a comprehensive review on heavy quarkonium physics [2]. It is certainly interesting to see whether the large discrepancy can be resolved by inclusion of NLO QCD corrections and relativistic corrections. Recently, in [23], the authors calculate the prompt $J/\psi + c\bar{c} + X$ production at NLO α_s including direct production and feeddown contribution mainly from the $\psi(2S)$, and find the NLO α_s corrections are large and positive, and could be helpful to settle the problem between experiment and theory. But we still need to know how large are the relativistic corrections, and whether they are positive or negative to the solution of the problem.

In this paper, we consider the relativistic corrections to both these exclusive and inclusive double-charm production processes based on NRQCD formulas in the color-singlet sector, since the color-octet contributions are negligible [21]. In order to avoid the model dependence in determining the long distance matrix elements, differing from [16], we determine the matrix elements of up to dimension-8 four fermion operators from the observed decay rates of J/ψ and η_c. We find that the relativistic effect on the double charmonium production $e^+e^- \to J/\psi + \eta_c$ is substantial and comparable to the estimate of [16]; whereas for the inclusive production $e^+e^- \to J/\psi + c\bar{c}$ the relativistic corrections are very small and negligible. The rest of this paper is organized as follows. In Section II, we give the general formulas of the production rates in NRQCD at v^2 order. Relativistic corrections to the exclusive process and the inclusive process are studied in Section III and Section IV respectively. A summary for the v^2 order corrections to these two processes will be given in Section V. In the Appendix we give some of the analytic results.

II. PRODUCTION CROSS SECTIONS IN NRQCD

In NRQCD the production and decay of charmonia are factorized into two parts, the short distance part that can be calculated perturbatively, and the long distance part can be estimated by lattice calculation, phenomenological models, or from other experimental observables. The long distance parts are related to the four fermion operators, characterized by the velocity v of the charm quark in the meson rest frame. The production cross sections of η_c and J/ψ up to v^2 order are [1]

$$\sigma(\eta_c) = \frac{F_1(^1S_0)}{m_c^2}\langle 0|\mathcal{O}_1(^1S_0^{\eta_c})|0\rangle + \frac{G_1(^1S_0)}{m_c^4}\langle 0|\mathcal{P}_1(^1S_0^{\eta_c})|0\rangle + O(v^4\sigma), \tag{3a}$$

$$\sigma(\psi) = \frac{F_1(^3S_1)}{m_c^2}\langle 0|\mathcal{O}_1(^3S_1^{\psi})|0\rangle + \frac{G_1(^3S_1)}{m_c^4}\langle 0|\mathcal{P}_1(^3S_1^{\psi})|0\rangle + O(v^4\sigma). \tag{3b}$$

The operators are defined as

$$\mathcal{O}_1^{\eta_c}(^1S_0) = \chi^\dagger\psi(a_{\eta_c}^\dagger a_{\eta_c})\psi^\dagger\chi, \tag{4a}$$

$$\mathcal{P}_1^{\eta_c}(^1S_0) = \frac{1}{2}[\chi^\dagger\psi(a_{\eta_c}^\dagger a_{\eta_c})\psi^\dagger(-\frac{i}{2}\overleftrightarrow{\mathbf{D}})^2\chi + \chi^\dagger(-\frac{i}{2}\overleftrightarrow{\mathbf{D}})^2\psi(a_{\eta_c}^\dagger a_{\eta_c})\psi^\dagger\chi], \tag{4b}$$

$$\mathcal{O}_1^{\psi}(^3S_1) = \chi^\dagger\sigma^i\psi(a_{\psi}^\dagger a_{\psi})\psi^\dagger\sigma^i\chi, \tag{4c}$$

$$\mathcal{P}_1^{\psi}(^3S_1) = \frac{1}{2}[\chi^\dagger\sigma^i\psi(a_{\psi}^\dagger a_{\psi})\psi^\dagger\sigma^i(-\frac{i}{2}\overleftrightarrow{\mathbf{D}})^2\chi + \chi^\dagger\sigma^i(-\frac{i}{2}\overleftrightarrow{\mathbf{D}})^2\psi(a_{\psi}^\dagger a_{\psi})\psi^\dagger\sigma^i\chi]. \tag{4d}$$

The short distance coefficients can be evaluated by the matching condition:

$$\sigma(Q\overline{Q})\Big|_{\text{pert QCD}} = \sum_n \frac{F_n(\Lambda)}{M^{d_n-4}}\langle 0|\mathcal{O}_n^{Q\overline{Q}}(\Lambda)|0\rangle\Big|_{\text{pert NRQCD}} \tag{5}$$

The left hand side of Eq. [5] can be calculated by the spinor projection method [24]. Furthermore (see, e.g.,[25][26]), the projection of $v(P/2-q)\overline{u}(P/2+q)$ onto a particular angular momentum state can be expressed in a Lorentz covariant form. The momenta of quark and antiquark in an arbitrary frame are respectively [27]:

$$\frac{1}{2}P + q = L(\frac{1}{2}P_r + \mathbf{q}), \tag{6a}$$

$$\frac{1}{2}P - q = L(\frac{1}{2}P_r - \mathbf{q}), \tag{6b}$$

where $P_r^\mu = (2E_q, \mathbf{0})$, $E_q = \sqrt{m^2+\mathbf{q}^2}$, and $2\mathbf{q}$ is the relative momentum between two quarks in the meson rest frame. L_μ^v is the boost tensor from the meson rest frame to an arbitrary frame.

In the meson rest frame the expression of projection onto a state of $S=0$ is

$$\sum_{\lambda_1\lambda_2} v(-\mathbf{q},\lambda_2)\overline{u}(\mathbf{q},\lambda_1)\langle\frac{1}{2},\lambda_1,\frac{1}{2},\lambda_2|0,0\rangle =$$
$$\frac{1}{\sqrt{2}}(E+m)(1-\frac{\boldsymbol{\alpha}\cdot\mathbf{q}}{E+m})\gamma_5\frac{1+\gamma_0}{2}(1+\frac{\boldsymbol{\alpha}\cdot\mathbf{q}}{E+m})\gamma_0. \tag{7}$$

In an arbitrary frame it becomes

$$\sum_{\lambda_1\lambda_2} v(q,\lambda_2)\overline{u}(q,\lambda_1)\langle\frac{1}{2},\lambda_1,\frac{1}{2},\lambda_2|0,0\rangle =$$
$$-\frac{1}{2\sqrt{2}(E+m)}(\frac{1}{2}\not{P}-\not{q}-m)\gamma_5\frac{\not{P}+2E}{2E}(\frac{1}{2}\not{P}+\not{q}+m). \tag{8}$$

Similarly, expressions of projection onto a state of $S=1$ in the rest frame of the meson and an arbitrary frame are:

$$\sum_{\lambda_1\lambda_2} v(-\mathbf{q},\lambda_2)\overline{u}(\mathbf{q},\lambda_1)\langle\frac{1}{2},\lambda_1;\frac{1}{2},\lambda_2|1,\epsilon\rangle =$$
$$\frac{1}{\sqrt{2}}(E+m)(1-\frac{\boldsymbol{\alpha}\cdot\mathbf{q}}{E+m})\boldsymbol{\alpha}\cdot\boldsymbol{\epsilon}\frac{1+\gamma_0}{2}(1+\frac{\boldsymbol{\alpha}\cdot\mathbf{q}}{E+m})\gamma_0, \tag{9a}$$

$$\sum_{\lambda_1\lambda_2} v(q,\lambda_2)\overline{u}(q,\lambda_1)\langle\frac{1}{2},\lambda_1;\frac{1}{2},\lambda_2|1,\epsilon\rangle =$$
$$-\frac{1}{2\sqrt{2}(E+m)}(\frac{1}{2}\not{P}-\not{q}-m)\not{\epsilon}\frac{\not{P}+2E}{2E}(\frac{1}{2}\not{P}+\not{q}+m). \tag{9b}$$

In our calculation the Dirac spinors are normalized as $\overline{u}u=-\overline{v}v=2m_c$. Then the short distance part of the cross section can be calculated at any order of v.

At order v^2 the cross section of $e^+e^-\to J/\psi+\eta_c$ and $e^+e^-\to J/\psi+c\overline{c}$ can be expressed as

$$\begin{aligned}\sigma(e^+e^-\to J/\psi+\eta_c) = \quad & \langle 0|\mathcal{O}_1(^1S_0^{\eta_c})|0\rangle\tfrac{\langle 0|\mathcal{O}_1(^3S_1^{\psi})|0\rangle}{3}\tfrac{1}{2s}\int\overline{M_0}\,d\,LIPS + \\ & \tfrac{\langle 0|\mathcal{P}_1(^3S_1^{\psi})|0\rangle}{3}\langle 0|\mathcal{O}_1(^1S_0^{\eta_c})|0\rangle\tfrac{1}{2s}\int\overline{M}_{1J/\psi}\,d\,LIPS + \\ & \langle 0|\mathcal{P}_1(^1S_0^{\eta_c})|0\rangle\tfrac{\langle 0|\mathcal{O}_1(^3S_1^{\psi})|0\rangle}{3}\tfrac{1}{2s}\int\overline{M}_{1\eta_c}\,d\,LIPS\,.\end{aligned} \tag{10a}$$

$$\begin{aligned}&\sigma(e^+e^-\to J/\psi+c\overline{c}) = \\ &\frac{\langle 0|\mathcal{O}_1(^3S_1^{\psi})|0\rangle}{3}\frac{1}{2s}\int\overline{N_0}\,d\,LIPS+\frac{\langle 0|\mathcal{P}_1(^3S_1^{\psi})|0\rangle}{3}\frac{1}{2s}\int\overline{N}_1\,d\,LIPS\,.\end{aligned} \tag{10b}$$

Here $LIPS$ means the Lorentz invariant phase space, bar means averaging spins over the initial states and summing spins over the final states, and $\overline{M_0},\overline{M}_{1J/\psi},\overline{M}_{1\eta_c},\overline{N_0},\overline{N_1}$ can be calculated perturbatively.



+2 flipped diagrams

FIG. 1: $e^+e^- \to (c\overline{c})_{^3S_1} + (c\overline{c})_{^1S_0}$

III. RELATIVISTIC CORRECTIONS TO $e^+e^- \to J/\psi + \eta_c$

A. Short distance part

There are four Feynman diagrams in the process $e^+e^- \to (c\overline{c})_{^3S_1}+(c\overline{c})_{^1S_0}$, shown in Fig.[1]. The amplitude of the process can be expanded in terms of the quark relative momentum in charmonium:

$$M(e^+e^- \to (c\overline{c})_{^3S_1} + (c\overline{c})_{^1S_0}) = (\frac{m_c}{E_1}\frac{m_c}{E_2})^{1/2} A(q_\psi, q_{\eta_c}) =$$
$$(\frac{m_c}{E_1}\frac{m_c}{E_2})^{1/2}(A(0,0) + q^\alpha_\psi \frac{\partial A}{\partial q^\alpha_\psi}\Big|_{q_\psi=q_{\eta c}=0} + q^\alpha_{\eta_c}\frac{\partial A}{\partial q^\alpha_{\eta_c}}\Big|_{q_\psi=q_{\eta c}=0} +$$
$$\frac{1}{2}q^\alpha_\psi q^\beta_\psi \frac{\partial^2 A}{\partial q^\alpha_\psi \partial q^\beta_\psi}\Big|_{q_\psi=q_{\eta c}=0} + \frac{1}{2}q^\alpha_{\eta_c} q^\beta_{\eta_c}\frac{\partial^2 A}{\partial q^\alpha_{\eta_c}\partial q^\beta_{\eta_c}}\Big|_{q_\psi=q_{\eta c}=0} + \dots), \tag{11}$$

and $A(q_\psi, q_{\eta_c})$ is expressed as

$$A(q_\psi, q_{\eta_c}) = \sum_{\lambda_1\lambda_2\lambda_3\lambda_4}\sum_{ijkl}\langle\frac{1}{2},\lambda_1;\frac{1}{2},\lambda_2|1,S_z\rangle\langle\frac{1}{2},\lambda_3;\frac{1}{2},\lambda_4|0,0\rangle\langle 3,i;\overline{3},j|1\rangle\langle 3,k;\overline{3},l|1\rangle$$
$$A(e^+e^- \to c_{\lambda 1,i}(\frac{P1}{2}+q_\psi)\overline{c}_{\lambda 2,j}(\frac{P1}{2}-q_\psi) + c_{\lambda 3,k}(\frac{P2}{2}+q_{\eta_c})\overline{c}_{\lambda 4,l}(\frac{P2}{2}-q_{\eta_c})), \tag{12}$$

where $\langle 3,i;\bar{3},j|1\rangle = \delta_{ij}/\sqrt{N_c}$ and $\langle 3,k;\bar{3},l|1\rangle = \delta_{kl}/\sqrt{N_c}$ are the color-SU(3), Clebsch-Gordon coefficients for $Q\bar{Q}$ pairs projecting onto a color singlet state. Using Eq.[8] and Eq.[9b], we can express $A(q_\psi, q_{\eta_c})$ in a covariant form. The factor $(\frac{m_c}{E_1}\frac{m_c}{E_2})^{1/2}$ comes from the relativistic normalization of the $c\overline{c}$ state, and $E_1 = \sqrt{m_c^2 + \mathbf{q}_\psi{}^2}$, $E_2 = \sqrt{m_c^2 + \mathbf{q}_{\eta c}{}^2}$.

For the S wave charmonium $q^\alpha q^\beta = \frac{1}{3}\mathbf{q}^2(-g^{\alpha\beta} + \frac{P^\alpha P^\beta}{P^2})$, where $P^2 = 4E^2, P\cdot q = 0$ and

the odd-power terms of v vanish. Then at leading order of v^2

$$|M|^2 = \frac{m_c}{E_1}\frac{m_c}{E_2}A(0,0)A^*(0,0) + \frac{1}{2}q_\psi^\alpha q_\psi^\beta A_\psi^{\alpha\beta}A^*(0,0) + \frac{1}{2}q_{\eta_c}^\alpha q_{\eta_c}^\beta A_{\eta_c}^{\alpha\beta}A^*(0,0) + \frac{1}{2}q_\psi^\alpha q_\psi^\beta A_\psi^{*\alpha\beta}A(0,0) + \frac{1}{2}q_{\eta_c}^\alpha q_{\eta_c}^\beta A_{\eta_c}^{*\alpha\beta}A(0,0), \tag{13}$$

where $A^{\alpha\beta} = \frac{\partial^2 A}{\partial q^\alpha \partial q^\beta}$, $A^{*\alpha\beta} = \frac{\partial^2 A^*}{\partial q^\alpha \partial q^\beta}$.

To obtain the short distance part $E = \sqrt{m_c^2 + \mathbf{q}^2}$ should also be expanded in $v^2 = \frac{\mathbf{q}^2}{m_c^2}$. Then the relation of $M_0, M_{1J/\psi}, M_{1\eta_c}$ to $|M|^2$ can be easily written down. Details of how to get the short distance coefficients from the covariant projection method can be found in e.g. [28]

$$\overline{M}_0 = \frac{1}{(2N_c m_c)^2}(A(0,0)A^*(0,0))\Big|_{\mathbf{q}^2_{J/\psi}=\mathbf{q}^2_{\eta c}=0}; \tag{14a}$$

$$\overline{M}_{1J/\psi} = \frac{1}{(2N_c m_c)^2}\Big(\frac{\partial(\frac{m_c^2}{E_1E_2}A(0,0)A^*(0,0))}{\partial(\mathbf{q}_\psi^2)}\Big|_{\mathbf{q}^2_{J/\psi}=\mathbf{q}^2_{\eta c}=0} + \frac{1}{6}(A_\psi^{\alpha\beta}\Pi_{\alpha\beta}(P_{J/\psi})A^*(0,0) + A_\psi^{*\alpha\beta}\Pi_{\alpha\beta}(P_{J/\psi})A(0,0))\Big)\Big|_{\mathbf{q}^2_{J/\psi}=\mathbf{q}^2_{\eta c}=0}; \tag{14b}$$

$$\overline{M}_{1\eta_c} = \frac{1}{(2N_c m_c)^2}\Big(\frac{\partial(\frac{m_c^2}{E_1E_2}A(0,0)A^*(0,0))}{\partial(\mathbf{q}_{\eta_c}^2)}\Big|_{\mathbf{q}^2_{J/\psi}=\mathbf{q}^2_{\eta c}=0} + \frac{1}{6}(A_{\eta_c}^{\alpha\beta}\Pi_{\alpha\beta}(P_{\eta_c})A^*(0,0) + A_{\eta_c}^{*\alpha\beta}\Pi_{\alpha\beta}(P_{\eta_c})A(0,0))\Big)\Big|_{\mathbf{q}^2_{J/\psi}=\mathbf{q}^2_{\eta c}=0}. \tag{14c}$$

Here $\overline{M}_0$ is just the leading order result, and it agrees with the previous result in ref.[7]

$$\overline{M}_0 = \frac{1}{9N_c^2}\frac{2048(s-16m_c^2)(1+\cos^2(\theta))(4\pi\alpha)^2(4\pi\alpha_s)^2e_q^2}{s^4}. \tag{15}$$

The expressions of $\overline{M}_{1J/\psi}$ and $\overline{M}_{1\eta_c}$ are

$$\overline{M}_{1J/\psi} = \frac{(1+\cos^2_\theta)(4\pi\alpha)^2(4\pi\alpha_s)^2e_q^2}{9N_c^2}\Big(\frac{512(16m_c^2-3s)}{m_c^2s^4} + \frac{512(2560m_c^4-592m_c^2s+27s^2)}{3m_c^2s^5}\Big), \tag{16a}$$

$$\overline{M}_{1\eta_c} = \frac{(1+\cos^2_\theta)(4\pi\alpha)^2(4\pi\alpha_s)^2e_q^2}{9N_c^2}\Big(\frac{-16384}{s^4} + \frac{1024(11s-80m_c^2)(s-16m_c^2)}{3m_c^2s^5}\Big), \tag{16b}$$

where $Nc = 3$, $e_q = \frac{2}{3}$, and θ is the angle between the J/ψ and the electron.

B. Long distance part

In this section we present a phenomenological estimate of the color-singlet production matrix elements which are extracted from charmonium decay data. It is known that up to errors of order v^4 the color-singlet production matrix elements are related to the decay matrix elements through vacuum saturation [1]. There are three independent NRQCD matrix elements at order v^2, $\langle 0|\mathcal{O}_1(^1S_0^{\eta_c})|0\rangle$, $\langle 0|\mathcal{O}_1(^3S_1^{\psi})|0\rangle$, and $\langle 0|\mathcal{P}_1(^1S_0^{\eta_c})|0\rangle = \langle 0|\mathcal{P}_1(^3S_1^{\psi})|0\rangle/3$ $(1+O(v^2))$. We can get them through the J/ψ leptonic decay $J/\psi \to e^+e^-$ and hadronic decay $J/\psi \to LH$, and the η_c photonic decay $\eta_c \to \gamma\gamma$. The theoretical results at next-to-leading order of α_s and v^2 for $J/\psi \to e^+e^-$, $J/\psi \to LH$ [a], and $\eta_c \to \gamma\gamma$ are summarized in ref.[25] as

$$\Gamma[\eta_c \to \gamma\gamma] = 2e_c^4\pi\alpha^2\Big((1-\frac{(20-\pi^2)\alpha_s}{3\pi})\frac{\langle 0|\mathcal{O}_1(^1S_0^{\eta_c})|0\rangle}{m_c^2} - \frac{4}{3}\frac{\langle 0|\mathcal{P}_1(^1S_0^{\eta_c})|0\rangle}{m_c^4}\Big), \tag{17a}$$

$$\Gamma[J/\psi \to e^+e^-] = \frac{2e_c^2\pi\alpha^2}{3}\Big((1-\frac{16\alpha_s}{3\pi})\frac{\langle 0|\mathcal{O}_1(^3S_1^{\psi})|0\rangle/3}{m_c^2} - \frac{4}{3}\frac{\langle 0|\mathcal{P}_1(^3S_1^{\psi})|0\rangle/3}{m_c^4}\Big), \tag{17b}$$

$$\Gamma[J/\psi \to LH] = (\frac{20\alpha_s^3}{243}(\pi^2-9))\Big((1-2.55\frac{\alpha_s}{\pi})\frac{\langle 0|\mathcal{O}_1(^3S_1^{\psi})|0\rangle/3}{m_c^2} - \frac{19\pi^2-132}{12\pi^2-108}\frac{\langle 0|\mathcal{P}_1(^3S_1^{\psi})|0\rangle/3}{m_c^4}\Big). \tag{17c}$$

Solving these equations at leading order of α_s (QCD radiative corrections not included), we get

$$\langle 0|\mathcal{O}_1(^1S_0^{\eta_c})|0\rangle = 0.286GeV^3, \langle 0|\mathcal{O}_1(^3S_1^{\psi})|0\rangle/3 = 0.295GeV^3, \tag{18a}$$

$$\frac{\langle 0|\mathcal{P}_1(^1S_0^{\eta_c})|0\rangle}{m_c^2} = \frac{\langle 0|\mathcal{P}_1(^3S_1^{\psi})|0\rangle}{3m_c^2} = 0.321\times 10^{-1}GeV^3, \tag{18b}$$

for m_c =1.5Gev and $\alpha_s = 0.26$. The experimental data of these decay rates can be found from [29], and we choose their central values $\Gamma[J/\psi \to e^+e^-] = 5.55$KeV, $\Gamma[J/\psi \to LH] =$ 69.3KeV, and $\Gamma[\eta_c \to \gamma\gamma] = 7.14$KeV. The matrix elements can be expressed as functions of the charm quark mass,

$$\langle 0|\mathcal{O}_1(^1S_0^{\eta_c})|0\rangle = 0.127m_c^2, \tag{19a}$$

$$\frac{\langle 0|\mathcal{O}_1(^3S_1^{\psi})|0\rangle}{3} = 0.131m_c^2, \tag{19b}$$

[a] We do not include the electromagnetic process of $J/\psi \to \gamma^* \to LH$.

$$\frac{\langle 0|\mathcal{P}_1(^1S_0^{\eta_c})|0\rangle}{m_c^2} = \frac{\langle 0|\mathcal{P}_1(^3S_1^{\psi})|0\rangle}{3m_c^2} = 0.014m_c^2 \tag{19c}$$

Including the QCD NLO radiative corrections, and doing the calculation in the same way as above, we have

$$\langle 0|\mathcal{O}_1(^1S_0^{\eta_c})|0\rangle = 0.432GeV^3, \langle 0|\mathcal{O}_1(^3S_1^{\psi})|0\rangle/3 = 0.573GeV^3, \tag{20a}$$

$$\frac{\langle 0|\mathcal{P}_1(^1S_0^{\eta_c})|0\rangle}{m_c^2} = \frac{\langle 0|\mathcal{P}_1(^3S_1^{\psi})|0\rangle}{3m_c^2} = 0.514 \times 10^{-1}GeV^3, \tag{20b}$$

for m_c =1.5Gev and $\alpha_s = 0.26$. And we then have

$$\langle 0|\mathcal{O}_1(^1S_0^{\eta_c})|0\rangle = 0.192m_c^2, \tag{21a}$$

$$\frac{\langle 0|\mathcal{O}_1(^3S_1^{\psi})|0\rangle}{3} = 0.255m_c^2, \tag{21b}$$

$$\frac{\langle 0|\mathcal{P}_1(^1S_0^{\eta_c})|0\rangle}{m_c^2} = \frac{\langle 0|\mathcal{P}_1(^3S_1^{\psi})|0\rangle}{3m_c^2} = 0.023m_c^2, \tag{21c}$$

as functions of m_c.

In [30], the authors express the next-to-leading order matrix elements in terms of the quark pole mass and the quarkonium mass, and in [16] the matrix elements are calculated based on the potential model. Differing from their methods, we get the production matrix elements by using the experimentally observed charmonium decay rates. As shown in[1], the difference between the color-singlet production and decay matrix elements are of order v^4. So our method should be valid at order v^2, and our estimates of the production matrix elements should be good numerically, if the high order QCD and v^2 corrections for the decays are small and the experimental errors are not large.

C. Numerical results and discussions

At $\sqrt{s}$ = 10.6GeV, with m_c = 1.5GeV and α_s = 0.26, using the matrix elements in Eqs.[18,19] determined from charmonium decays with v^2 corrections but without QCD radiative corrections, and making the phase space integral, the leading order cross section of $e^+e^- \to J/\psi + \eta_c$ (LO means for the short-distance part, since for the long-distance matrix elements the v^2 corrections are already included) is 3.07 fb. The relativistic correction



FIG. 2: $e^+e^- \to J/\psi + \eta_c$ cross sections with relativistic corrections to long-distance matrix elements extracted from charmonium decays (without NLO QCD radiative corrections). The lower line represents the LO result in v, and the upper line represents the result with v^2 corrections to the short-distance coefficients. Here the coupling constant is fixed as $\alpha_s = 0.26$.

contributes 0.798 fb, which gives about 26% enhancement and results in 3.87 fb for the cross section. The cross sections as functions of m_c are shown in Fig[2]. The lower line represents the LO result in v, and the upper line represents the result with v^2 corrections. Since the long-distance matrix elements are proportional to the squared quark mass, and the short distance coefficients are found to be quite stable when m_c changes, so the cross section goes down as m_c becomes smaller. However, these results are obtained for fixed value of $\alpha_s = 0.26$, and if the running of $\alpha_s = \alpha_s(2m_c)$ is assumed, the m_c dependence of cross sections will be substantially changed, making the cross section at m_c=1.4 GeV close to that at m_c=1.6 GeV.

If using the matrix elements in Eqs.[20,21] determined from charmonium decays with both v^2 and α_s corrections, at order v^0 the cross section is 9.02fb, and at order v^2 the cross section is 11.26fb for $\alpha_s = 0.26$ and $m_c = 1.5$Gev. When m_c varies from 1.4Gev to 1.6Gev, the cross section goes up from 9.18fb to 13.43fb.

The experiment result in Eq.[1] is about an order of magnitude larger than the leading



FIG. 3: $e^+e^- \to J/\psi + \eta_c$ cross sections with relativistic corrections to long-distance matrix elements extracted from charmonium decays (with NLO QCD radiative corrections). The lower line represents the LO result in v, and the upper line represents the result with v^2 corrections to the short-distance coefficients. Here the coupling constant is fixed as $\alpha_s = 0.26$. Note that the QCD radiative corrections to the short-distance coefficients (with K=1.8) are included for the upper line but not the lower line.

order result[6, 7, 8]. The result at order v^2 shows that relativistic corrections can enhance both the short distance coefficients and the long distance matrix elements. The next-to-leading order v^2 coefficient $(\overline{M}_{1J/\psi} + \overline{M}_{1\eta_c})m_c^2$ is about 2.2 times larger than $\overline{M}$ for $m_c =$ 1.5Gev and $\alpha_s = 0.26$, and the matrix elements of $\langle 0|\mathcal{O}_1(^1S_0^{\eta_c})|0\rangle$ and $\langle 0|\mathcal{O}_1(^3S_1^{\psi})|0\rangle$ also become about 1.17 times larger than that in the leading order calculation. These make the cross section become 1.7 times larger after including the relativistic effect. If we determine the matrix elements including the QCD radiative correction, at v^2 order the theoretical result is 11.26fb for $\alpha_s = 0.26$ and $m_c = 1.5$Gev. It shows that the relativistic corrections are significant.

In the above discussions, we have not considered the QCD radiative corrections to the short-distance coefficients as shown in [9]. If we further combine the NLO QCD corrections [9] with the v^2 corrections with a fixed value $\alpha_s = 0.26$, then the cross section will go from

TABLE I: Experimental and calculated cross sections of $\sigma[e^+e^- \to J/\psi + \eta_c]$ with m_c =1.5 GeV and α_s=0.26. See text for the definitions of $\sigma_{LO(\alpha_s,v^2)}$, $\sigma_{NLO(\alpha_s)}$, $\sigma_{NLO(v^2)}$, and $\sigma_{NLO(\alpha_s,v^2)}$.

Experimental Result					
$\sigma_{Belle}[e^+e^- \to J/\psi + \eta_c] \times \mathcal{B}^{\eta_c}[\geq 2]$(fb)			$\sigma_{Babar}[e^+e^- \to J/\psi + \eta_c] \times \mathcal{B}^{\eta_c}[\geq 2]$(fb)		
$25.6 \pm 2.8 \pm 3.4$			$17.6 \pm 2.8 \pm 2.1$		
Theoretical Result					
$\langle 0\|\mathcal{O}_1(^1S_0^{\eta_c})\|0\rangle$	$\frac{\langle 0\|\mathcal{O}_1(^3S_1^{\psi})\|0\rangle}{3}$	$\frac{\langle 0\|\mathcal{P}_1(^1S_0^{\eta_c})\|0\rangle}{m_c^2}$	$\frac{\langle 0\|\mathcal{P}_1(^3S_1^{\psi})\|0\rangle}{3m_c^2}$	α_s	σ (fb)
0.243Gev^3	0.252Gev^3	0	0	$\alpha_s = 0.26$	$\sigma_{LO(\alpha_s,v^2)} = 2.26$
0.337Gev^3	0.450Gev^3	0	0	$\alpha_s = 0.26$	$\sigma_{NLO(\alpha_s)} = 10.92$
0.286Gev^3	0.295Gev^3	0.0321Gev^3	0.0321Gev^3	$\alpha_s = 0.26$	$\sigma_{NLO(v^2)} = 3.87$
0.432Gev^3	0.573Gev^3	0.0514Gev^3	0.0514Gev^3	$\alpha_s = 0.26$	$\sigma_{NLO(\alpha_s,v^2)} = 20.04$

16.8fb to 23.0fb when m_c varies from 1.4GeV to 1.6 GeV, as shown in Fig[3]. The calculated cross sections of $\sigma[e^+e^- \to J/\psi + \eta_c]$ are listed in Table I, where $\sigma_{LO(\alpha_s,v^2)}$ means the cross section at leading order in both α_s and v^2; $\sigma_{NLO(\alpha_s)}$ means that obtained by using the short distance part with α_s corrections [9], and the long-distance matrix elements also with α_s corrections extracted from both $J/\psi \to e^+e^-$ and $\eta_c \to 2\gamma$ (not from $J/\psi \to e^+e^-$ alone as was extracted in [9]) with all at leading order in v^2; $\sigma_{NLO(v^2)}$ means that obtained with v^2 corrections but without α_s corrections; and $\sigma_{NLO(\alpha_s,v^2)}$ means that obtained by combining both α_s and v^2 corrections, where the matrix element are taken from Eq.[20].

We see that with both QCD radiative corrections and relativistic corrections, the discrepancy between experiment and theory for $e^+e^- \to J/\psi + \eta_c$ could be largely resolved. Our result is consistent with [16]. In our approach the long-distance matrix elements are extracted from experimental data of J/ψ and η_c decays, and therefore are model-independent. The main uncertainties may come from the higher order corrections and the errors in the measurements. The relativistic effects on the double charmonium production estimated in our approach are milder than some results obtained by using the light-cone methods.



FIG. 4: Feynman diagrams for $e^+ + e^- \to \gamma^* \to J/\psi + c\bar{c}$.

IV. RELATIVISTIC CORRECTIONS TO $e^+e^- \to J/\psi + c\overline{c}$

The long distance matrix elements in process $e^+e^- \to J/\psi + c\overline{c}$ are the same as in process $e^+e^- \to J/\psi + \eta_c$, so we just use the result above, and only give the detailed calculation for the short distance part.

A. Short distance part

There are four Feynman diagrams in the process $e^+e^- \to (C\overline{C})_{^3S_1} + C\overline{C}$, which are shown in Fig[4]. As in the last section the amplitude can be expanded in terms of the quark relative momentum and reads

$$M(e^+e^- \to (C\overline{C})_{^3S_1} + C\overline{C}) = (\frac{m_c}{E})^{1/2} A(q_\psi) =$$
$$(\frac{m_c}{E})^{1/2}(A(0) + q_\psi^\alpha \frac{\partial A}{\partial q_\psi^\alpha}\Big|_{q_\psi=0} + \frac{1}{2} q_\psi^\alpha q_\psi^\beta \frac{\partial^2 A}{\partial q_\psi^\alpha \partial q_\psi^\beta}\Big|_{q_\psi=0} + \dots), \tag{22}$$

where

$$A(q_\psi) = \sum_{\lambda_1\lambda_2}\sum_{ij} \langle \frac{1}{2}, \lambda_1, \frac{1}{2}, \lambda_2 | 1, S_z\rangle \langle 3, i; \overline{3}, j | 1\rangle A(e^+e^- \to C_{\lambda 1, i}(\frac{P}{2} + q_\psi)\overline{C}_{\lambda 2, j}(\frac{P}{2} - q_\psi) + C\overline{C}) \tag{23}$$

At leading order of v^2

$$|M|^2 = \frac{m_c}{E_1} A(0)A^*(0) + \frac{1}{2} q_\psi^\alpha q_\psi^\beta A_{\alpha\beta} A^*(0) + \frac{1}{2} q_\psi^\alpha q_\psi^\beta A^*_{\alpha\beta} A(0), \tag{24}$$

where $A_{\alpha\beta} = \frac{\partial^2 A}{\partial q^\alpha \partial q^\beta}$, and $A^*_{\alpha\beta} = \frac{\partial^2 A^*}{\partial q^\alpha \partial q^\beta}$. Then

$$\overline{N}_0 = \frac{1}{2N_c m_c}(A(0)A^*(0))\Big|_{\mathbf{q}^2_{J/\psi}=0}, \tag{25a}$$

$$\overline{N}_1 = \frac{1}{2N_c m_c}(\frac{\partial(\frac{m_c}{E}A(0)A^*(0))}{\partial(\mathbf{q}^2_\psi)}\Big|_{\mathbf{q}^2_{J/\psi}=0}$$
$$+\frac{1}{6}(A^{\alpha\beta}_\psi \Pi_{\alpha\beta}(P_{J/\psi})A^*(0) + A^{*\alpha\beta}_\psi \Pi_{\alpha\beta}(P_{J/\psi})A(0))\Big|_{\mathbf{q}^2_{J/\psi}=0}). \tag{25b}$$

The expressions of $A(0)A^*(0)$ and $\Pi^{\alpha\beta}(A_{\alpha\beta}A^*(0) + A^*_{\alpha\beta}A(0))$ can be written in terms of some dimensionless variables $z_i = 2E_i/\sqrt{s}$, $\overrightarrow{q_i} = 2\overrightarrow{p_i}/\sqrt{s}$, $x_i = \cos\theta_i$ and $\delta_i = 2m_i/\sqrt{s}$. Here $\sqrt{s}$ is the total energy in the center of mass frame, p^μ_3, p^μ_4, p^μ_5 are the four-momenta of the final state J/ψ, charm quark and anticharm quark respectively, $m_i^2 = p_i^2$, and θ_i is the angle between state i and the electron. The scalar productions are expressed as follows:

$$\begin{aligned}
&p_1.p_3 = \frac{s}{4}(z_3 - q_3 x_3);\ p_2.p_3 = \frac{s}{4}(z_3 + q_3 x_3);\ p_4.p_5 = \frac{s}{8}(4 - 4z_3 + \delta_3^2 - \delta_4^2 - \delta_5^2);\\
&p_1.p_4 = \frac{s}{4}(z_4 - (q_- x_- - q_3 x_3)/2);\ p_2.p_4 = \frac{s}{4}(z_4 + (q_- x_- - q_3 x_3)/2);\\
&p_1.p_5 = \frac{s}{4}(z_5 + (q_- x_- + q_3 x_3)/2);\ p_2.p_5 = \frac{s}{4}(z_5 - (q_- x_- + q_3 x_3)/2);\\
&p_3.p_4 = \frac{s}{8}(4 - \delta_3^2 - 4z_5);\ p_3.p_5 = \frac{s}{8}(4 - \delta_3^2 - 4z_4);\ p_1.p_2 = \frac{s}{2};
\end{aligned} \tag{26}$$

where $z_- = z_4 - z_5$, $q_- = |\overrightarrow{q_4} - \overrightarrow{q_5}| = \sqrt{4 - 4z_3 + z_-^2 + \delta_3^2 - 2\delta_4^2 - 2\delta_5^2}$, $q_3 = |\overrightarrow{q_3}| = \sqrt{z_3 - \delta_3^2}$, $x_- = \cos\theta_-$, and θ_- is the angle between $\overrightarrow{q_-}$ and the electron.

B. Numerical result

In principle, the phase space calculation also contains relativistic corrections (e.g., by inclusion of the binding energy in a meson). But for simplicity, we calculate the phase space integration by assuming $m_{J/\psi} = 2m_c$ (the effect due to this simplification is very small). Then the three-body final state phase space expressed in terms of the variables defined above becomes

$$\begin{aligned}
d\Phi_3 &= (2\pi)^4\ \delta^4(p1 + p2 - p3 - p4 - p5)\prod_{i=3}^{5}\frac{d^3 p_i}{2E_i}\\
&= \frac{s}{32(2\pi)^4}\frac{dz_3 dx_3 dz_- dw}{\sqrt{(1-K^2)(1-x_3^2) - w^2}},
\end{aligned} \tag{27}$$

where

$$K = \frac{z_-(2-z_3)}{q_3 q_-}, \tag{28a}$$

$$w = x_- + K x_3. \tag{28b}$$

The limits of those variables are

$$\delta \le z_3 \le 1, \tag{29a}$$

$$-1 \le x_3 \le 1, \tag{29b}$$

$$-\sqrt{\frac{(z_3^2-\delta^2)(4-4z_3)}{4+\delta^2-4z_3}} \le z_- \le \sqrt{\frac{(z_3^2-\delta^2)(4-4z_3)}{4+\delta^2-4z_3}}, \tag{29c}$$

$$-\sqrt{(1-K^2)(1-x_3^2)} \le w \le \sqrt{(1-K^2)(1-x_3^2)}, \tag{29d}$$

where $\delta = 4m_c/\sqrt{s}$.

The expressions of M_0 and M_1 are lengthy, so we only give the expression for the differential cross section. The differential cross section of unpolarized J/ψ production in association with $c\bar{c}$ in e^+e^- annihilation can be expressed as

$$\frac{d^2\sigma}{dE_3 d\cos\theta_3}(e^+e^- \to \gamma^* \to \psi + X) = S(E_3)(1+\alpha\cos^2\theta_3), \tag{30}$$

where E_3 is the energy of J/ψ, and θ_3 is the angle between J/ψ and the electron. When expressing the above equation by using the new parameters it becomes

$$\frac{d^2\sigma}{dz_3 dx_3}(e^+e^- \to \gamma^* \to \psi + X) = S(z_3)(1+\alpha(z_3)x_3^2). \tag{31}$$

The leading order result of $S(z_3)$ and $\alpha(z_3)$ are in agreement with those in [17, 21]. We give the next-to-leading order result of $S(z_3)$ and $\alpha(z_3)$ in the Appendix **A**. Using the long-distance matrix elements in Eq.[18] (without NLO α_s corrections in the charmonium decay widths), the leading order cross section of $e^+ + e^- \to J/\psi + c\overline{c}$ with $m_c = 1.5$ Gev, $\sqrt{s} = 10.6$ Gev, $\alpha_s = 0.26$ is estimated to be 110 fb,[b] and the next-to-leading order v^2 correction is only 0.42 fb, which gives about less than a half percent enhancement. As a result, the relativistic correction to $e^+ + e^- \to J/\psi + c\overline{c}$ is found to be very small and negligible, in contrast to the exclusive double charmonium production process $e^+ + e^- \to J/\psi + \eta_c$, in which the relativistic correction is very significant. In fact, we find that in $e^+ + e^- \to J/\psi + c\overline{c}$ the

[b] This value is smaller than 148 fb given in [21], because here a smaller value of the wave function squared at the origin is extracted from the J/ψ data than that used in [21] from potential model calculations.



FIG. 5: R_v as a function of $\sqrt{s}$. Here R_v is the ratio of the correction at order v^2 to the leading order result for the cross section of $e^+ + e^- \to J/\psi + c\overline{c}$.

short distance part $\bar{N}_1 m_c^2$ at order v^2 is much smaller than the leading order term $\bar{N}_0$, i.e., $\frac{\bar{N}_1 m_c^2}{\bar{N}_0} \ll 1$. Here $\bar{N}_1$ in Eq.[25b] is the sum of two terms, and they both are small and have different signs, so their sum becomes even tiny. So, the tiny effect of relativistic corrections on the rate of $e^+ + e^- \to J/\psi + c\overline{c}$ is due to the smallness of the short-distance coefficient correction, regardless of the long-distance matrix elements.

When $\sqrt{s}$ is larger, the ratio of the correction at order v^2 to the leading order contribution even changes sign from positive to negative values, but its value is always small. This is shown in Fig[5], where the values of the parameters are the same as used above.

If we use the enhanced matrix elements in Eq.[20], which are obtained by including NLO α_s corrections in the charmonium decay widths, for $m_c = 1.5$Gev, $\sqrt{s} = 10.6$Gev, $\alpha_s = 0.26$, the leading order result is about 214fb, and the relativistic correction at order v^2 is only 0.67fb.

C. Discussions

Since for $e^+ + e^- \to J/\psi + c\overline{c}$ the conflict between experiment and theory at leading order in both α_s and v^2 is quite serious, a number of attempts have been made to solve this problem. In [22] the authors considered the production of $J/\psi + c\overline{c}$ through two photons, and found that the contribution of two photon process may be comparable to that of one photon process when $\sqrt{s}$ is large, say, larger than $\sqrt{s} = 20$Gev, but at $\sqrt{s} = 10.6$Gev, it is

only 15% of the one photon contribution. Authors in [31] considered the process when the mass of the charm quark can be neglected. In [32] the result in factorization method was compared with the duality method. Other suggestions can also be found in e.g. [33, 34]. Despite of these efforts, a satisfactory resolution is still needed. Recently, the authors of ref.[23] have calculated the next-to-leading order QCD corrections to the direct $J/\psi+c\bar{c}+X$ production in $e^+ + e^-$ annihilation at $\sqrt{s} = 10.6$GeV, and with $m_c = 1.5$GeV and $\alpha_s = 0.26$ their result is

$$\sigma_{direct}(e^+e^- \to J/\psi + c\overline{c}) \approx 0.33\text{pb}, \tag{32}$$

It enhances the leading order result by a factor of 1.8. Further including the contributions from $\psi(2S)$ and other states, the cross section of prompt $J/\psi + c\overline{c}$ production becomes

$$\sigma_{prompt}(e^+e^- \to J/\psi + c\overline{c} + X) \approx 0.51\text{pb}. \tag{33}$$

It is about 60% of the Belle value in Eq.[2]. This result shows that the QCD radiative corrections to $e^+ + e^- \to J/\psi + c\overline{c}$ is essential.

Here, when we further consider the relativistic corrections at order v^2, we find that in contrast to the $J/\psi + \eta_c$ production, the relativistic corrections to $J/\psi + c\overline{c}$ production are very small. The relativistic corrections can only improve the leading order result by enlarging the long distance matrix elements, and have little effect on the short distance part. As a result, the relativistic corrections to $e^+ + e^- \to J/\psi + c\overline{c}$ are tiny and negligible.

The physical reason for the difference between exclusive and inclusive processes largely lies in the fact that in the exclusive process $e^+e^- \to J/\psi + \eta_c$ the virtuality of the gluon that converts into a charm quark pair takes its maximum value $\frac{s}{4}$ in the nonrelativistic limit (with zero relative momentum between quarks in charmonium), and introducing the relative momentum can substantially reduce the gluon virtuality, and hence enhance the short-distance coefficient and the cross section; whereas for the inclusive process $e^+ + e^- \to J/\psi + c\overline{c}$, the charm quark fragmentation $c \to J/\psi + c$ ($\bar{c} \to J/\psi + \bar{c}$) is significant, in which the quark relative momentum has little effect on the virtuality of the gluon. Therefore, our result for the next-to-leading v^2 corrections may indicate that the relativistic correction is not a good direction to solve the problem of discrepancy between theory and experiment in $J/\psi + c\bar{c}$ production. Further studies for QCD radiative corrections and other possible new mechanisms related to the double charm production could be more useful.

V. SUMMARY

In this paper, we have studied the relativistic corrections at order v^2 to the double-charm production processes $e^+e^- \to J/\psi + \eta_c$ and $e^+e^- \to J/\psi + c\overline{c}$ at B factories in the framework of non-relativistic quantum chromodynamics. The short-distance parts of production cross sections are calculated perturbatively, while the long-distance matrix elements are extracted from experimental data for J/ψ and η_c decays up to errors of order v^4, and therefore are model-independent. The main uncertainties may come from the higher order corrections and the errors in the measurements. Our results show that the relativistic correction to the exclusive process $e^+e^- \to J/\psi + \eta_c$ is significant, which, when combined together with the next-to-leading order α_s corrections, could resolve the large discrepancy between theory and experiment. It can be clearly seen from TABLE I that for the cross section the relativistic correction alone gives an enhancement factor of 1.7 while the combination of relativistic correction with QCD radiative correction results in a much larger enhancement factor of 9. This conclusion is consistent with [16], in which, however, the long-distance matrix elements are estimated by using potential model calculations. The relativistic effects on the $J/\psi\eta_c$ production estimated in our approach are milder than some results obtained by using the light-cone methods. On the other hand, for the inclusive process $e^+e^- \to J/\psi + c\overline{c}$ we find that the relativistic correction is tiny and negligible, and therefore not helpful in resolving the discrepancy between theory and experiment. The physical reason for the above difference between exclusive and inclusive processes largely lies in the fact that in the exclusive process the relative momentum between quarks in charmonium substantially reduces the virtuality of the gluon that converts into a charm quark pair, but this is not the case for the inclusive process, in which the charm quark fragmentation $c \to J/\psi + c$ ($\bar{c} \to J/\psi + \bar{c}$) is significant, and QCD radiative corrections can be more essential. Further studies are needed to understand the large ratio of the cross section of $e^+e^- \to J/\psi + c\overline{c}$ to the cross section of $e^+e^- \to J/\psi + anything$, which is measured to be in the range of 0.6-0.8 by Belle.

Acknowledgments

We thank G.T. Bodwin for useful comments. This work was supported in part by the National Natural Science Foundation of China (No 10421503, No 10675003), the Key Grant Project of Chinese Ministry of Education (No 305001), and the Research Found for Doctorial Program of Higher Education of China.

APPENDIX A

The formulas of $S(z_3)$ and $\alpha_(z_3)$ are given as follows:

$$S(z_3) = \frac{2e_c^2\pi\alpha^2\alpha_s^2}{3^5 s^3\delta^4 z_3^4\sqrt{1-z_3}(2-z_3)^6(z_3^2-\delta^2)}\frac{\langle 0|\mathcal{P}_1(^3S_1^{\psi})|0\rangle}{m_c}\times$$

$$\Big(-(98304\,{z_3}^5+98304\,{z_3}^6+696320\,{z_3}^7-6963200\,{z_3}^8+14895104\,{z_3}^9-13801472\,{z_3}^{10}+$$

$$6104576\,{z_3}^{11}-1221632\,{z_3}^{12}+93696\,{z_3}^{13}-737280\,{z_3}^3\,\delta^2+3047424\,{z_3}^4\,\delta^2-6987776\,{z_3}^5\,\delta^2+$$

$$15003648\,{z_3}^6\,\delta^2-22528000\,{z_3}^7\,\delta^2+17085440\,{z_3}^8\,\delta^2-4745728\,{z_3}^9\,\delta^2-411904\,{z_3}^{10}\,\delta^2+$$

$$299584\,{z_3}^{11}\,\delta^2-26176\,{z_3}^{12}\,\delta^2+737280\,z_3\,\delta^4-4030464\,{z_3}^2\,\delta^4+7950336\,{z_3}^3\,\delta^4-$$

$$7868416\,{z_3}^4\,\delta^4+3901440\,{z_3}^5\,\delta^4+2526208\,{z_3}^6\,\delta^4-6324224\,{z_3}^7\,\delta^4+3657984\,{z_3}^8\,\delta^4-$$

$$540096\,{z_3}^9\,\delta^4-42816\,{z_3}^{10}\,\delta^4+3632\,{z_3}^{11}\,\delta^4+442368\,\delta^6-1124352\,z_3\,\delta^6+387072\,{z_3}^2\,\delta^6+$$

$$948224\,{z_3}^3\,\delta^6-1466368\,{z_3}^4\,\delta^6+1291008\,{z_3}^5\,\delta^6-68864\,{z_3}^6\,\delta^6-493440\,{z_3}^7\,\delta^6+99136\,{z_3}^8\,\delta^6+$$

$$34552\,{z_3}^9\,\delta^6-664\,{z_3}^{10}\,\delta^6+211968\,\delta^8-489984\,z_3\,\delta^8+357888\,{z_3}^2\,\delta^8-14336\,{z_3}^3\,\delta^8-$$

$$365056\,{z_3}^4\,\delta^8+357696\,{z_3}^5\,\delta^8-42464\,{z_3}^6\,\delta^8-23104\,{z_3}^7\,\delta^8-7356\,{z_3}^8\,\delta^8-138\,{z_3}^9\,\delta^8+$$

$$39168\,\delta^{10}-93696\,z_3\,\delta^{10}+95232\,{z_3}^2\,\delta^{10}-20224\,{z_3}^3\,\delta^{10}-52576\,{z_3}^4\,\delta^{10}+23392\,{z_3}^5\,\delta^{10}+$$

$$3904\,{z_3}^6\,\delta^{10}+432\,{z_3}^7\,\delta^{10}+45\,{z_3}^8\,\delta^{10}+3456\,\delta^{12}-9216\,z_3\,\delta^{12}+8352\,{z_3}^2\,\delta^{12}+$$

$$384\,{z_3}^3\,\delta^{12}-2520\,{z_3}^4\,\delta^{12}+192\,{z_3}^5\,\delta^{12}-90\,{z_3}^6\,\delta^{12})\frac{4z_3(1-z_3)}{3(2-z_3)^2}\sqrt{\frac{z_3^2-\delta^2}{\delta^2-4z_3+4}}$$

$$-192(256-512\,z_3+320\,{z_3}^2-64\,{z_3}^3+16\,z_3^4-16\,z_3^5+4\,z_3^6-128\,\delta^2+64\,z_3\,\delta^2+$$

$$240\,z_3^2\,\delta^2-224\,z_3^3\,\delta^2+20\,z_3^4\,\delta^2+16\,z_3^5\,\delta^2+48\,\delta^4-96\,z_3\,\delta^4+24\,z_3^2\,\delta^4+24\,z_3^3\,\delta^4+3\,z_3^4\,\delta^4)$$

$$\delta^4 z_3^4(1-z_3)\arctan\frac{z_3^2-\delta^2}{\delta^2-4z_3+4}$$

$$(8192\,{z_3}^6-102400\,z_3^7+262144\,z_3^8-282624\,z_3^9+146944\,z_3^{10}-35584\,z_3^{11}+3072\,z_3^{12}+$$

$$69632\,z_3^3\,\delta^2-299008\,z_3^4\,\delta^2+542720\,z_3^5\,\delta^2-564224\,z_3^6\,\delta^2+331264\,z_3^7\,\delta^2-87808\,z_3^8\,\delta^2+$$

$$1664\,z_3^9\,\delta^2+1472\,z_3^{10}\,\delta^2+336\,z_3^{11}\,\delta^2-61440\,z_3\,\delta^4+299008\,z_3^2\,\delta^4-486400\,z_3^3\,\delta^4+$$

$$388096\,z_3^4\,\delta^4-180480\,z_3^5\,\delta^4+65280\,z_3^6\,\delta^4-19008\,z_3^7\,\delta^4-2880\,z_3^8\,\delta^4+3264\,z_3^9\,\delta^4-$$

$$288\,z_3^{10}\,\delta^4-36864\,\delta^6+72192\,z_3\,\delta^6-16896\,z_3^2\,\delta^6-64256\,z_3^3\,\delta^6+65664\,z_3^4\,\delta^6-$$

$$15360\,z_3^5\,\delta^6-6240\,z_3^6\,\delta^6+304\,z_3^7\,\delta^6+2216\,z_3^8\,\delta^6+6\,z_3^9\,\delta^6-8448\,\delta^8+14336\,z_3\,\delta^8-$$

$$6144\,z_3^2\,\delta^8-4864\,z_3^3\,\delta^8+1440\,z_3^4\,\delta^8+2624\,z_3^5\,\delta^8-1056\,z_3^6\,\delta^8+15\,z_3^8\,\delta^8-$$

$$1152\,\delta^{10}+3072\,z_3\,\delta^{10}-2592\,z_3^2\,\delta^{10}+1344\,z_3^3\,\delta^{10}-696\,z_3^4\,\delta^{10}+240\,z_3^5\,\delta^{10}-30\,z_3^6\,\delta^{10})$$

$$\delta^2\sqrt{1-z_3}\ln\frac{z_3\sqrt{\delta^2-4z_3+4}+2\sqrt{(1-z_3)(z_3^2-\delta^2)}}{z_3\sqrt{\delta^2-4z_3+4}-2\sqrt{(1-z_3)(z_3^2-\delta^2)}}\Big), \qquad \text{(A1)}$$

$$\alpha(z_3)S(z_3)=\frac{2e_c^2\pi\alpha^2\alpha_s^2}{3^5s^3\delta^4z_3^4\sqrt{1-z_3}(2-z_3)^6(z_3^2-\delta^2)}\frac{\langle 0|\mathcal{P}_1(^3S_1^{\psi})|0\rangle}{m_c}\times$$
$$\Big(-(98304\,{z_3}^7+98304\,z_3^8+696320\,z_3^9-6963200\,z_3^{10}+14895104\,z_3^{11}-$$
$$13801472\,z_3^{12}+6104576\,z_3^{13}-1221632\,z_3^{14}+93696\,z_3^{15}-835584\,z_3^5\,\delta^2+3342336\,z_3^6\,\delta^2-$$
$$11059200\,z_3^7\,\delta^2+31518720\,z_3^8\,\delta^2-49162240\,z_3^9\,\delta^2+37194752\,z_3^{10}\,\delta^2-11941888\,z_3^{11}\,\delta^2+$$
$$816896\,z_3^{12}\,\delta^2+153152\,z_3^{13}\,\delta^2-26176\,z_3^{14}\,\delta^2-1867776\,z_3^3\,\delta^4+6881280\,z_3^4\,\delta^4-$$
$$4657152\,z_3^5\,\delta^4-19660800\,z_3^6\,\delta^4+43751424\,z_3^7\,\delta^4-31881216\,z_3^8\,\delta^4+6889984\,z_3^9\,\delta^4+$$
$$366080\,z_3^{10}\,\delta^4+119808\,z_3^{11}\,\delta^4+31744\,z_3^{12}\,\delta^4-976\,z_3^{13}\,\delta^4+245760\,z_3\,\delta^6-1622016\,z_3^2\,\delta^6+$$
$$3229696\,z_3^3\,\delta^6-1411072\,z_3^4\,\delta^6+2289664\,z_3^5\,\delta^6-12958720\,z_3^6\,\delta^6+16642816\,z_3^7\,\delta^6-$$
$$7645696\,z_3^8\,\delta^6+1814848\,z_3^9\,\delta^6-524672\,z_3^{10}\,\delta^6+16520\,z_3^{11}\,\delta^6-1240\,z_3^{12}\,\delta^6+147456\,\delta^8-$$
$$522240\,z_3\,\delta^8+967680\,z_3^2\,\delta^8-2955776\,z_3^3\,\delta^8+6496768\,z_3^4\,\delta^8-5705984\,z_3^5\,\delta^8+1570816\,z_3^6\,\delta^8-$$
$$345536\,z_3^7\,\delta^8+177184\,z_3^8\,\delta^8+68040\,z_3^9\,\delta^8+3420\,z_3^{10}\,\delta^8+294\,z_3^{11}\,\delta^8+46080\,\delta^{10}+10752\,z_3\,\delta^{10}-$$
$$254208\,z_3^2\,\delta^{10}+92672\,z_3^3\,\delta^{10}+144640\,z_3^4\,\delta^{10}+86976\,z_3^5\,\delta^{10}+9920\,z_3^6\,\delta^{10}-81632\,z_3^7\,\delta^{10}-$$
$$996\,z_3^8\,\delta^{10}-2070\,z_3^9\,\delta^{10}+45\,z_3^{10}\,\delta^{10}+16128\,\delta^{12}-18432\,z_3\,\delta^{12}-15744\,z_3^2\,\delta^{12}+65408\,z_3^3\,\delta^{12}-$$
$$116416\,z_3^4\,\delta^{12}+61312\,z_3^5\,\delta^{12}-4136\,z_3^6\,\delta^{12}+1800\,z_3^7\,\delta^{12}+45\,z_3^8\,\delta^{12}+3456\,\delta^{14}-9216\,z_3\,\delta^{14}+$$
$$8352\,z_3^2\,\delta^{14}+384\,z_3^3\,\delta^{14}-2520\,z_3^4\,\delta^{14}+192\,z_3^5\,\delta^{14}-90\,z_3^6\,\delta^{14})\frac{4z_3(1-z_3)}{3(2-z_3)^2}\sqrt{\frac{z_3^2-\delta^2}{\delta^2-4z_3+4}}$$
$$+192(256-512\,z_3+320\,z_3^2-64\,z_3^3+16\,z_3^4-16\,z_3^5+4\,z_3^6-128\,\delta^2+64\,z_3\,\delta^2+240\,z_3^2\,\delta^2-$$
$$224\,z_3^3\,\delta^2+20\,z_3^4\,\delta^2+16\,z_3^5\,\delta^2+48\,\delta^4-96\,z_3\,\delta^4+24\,z_3^2\,\delta^4+24\,z_3^3\,\delta^4+3\,z_3^4\,\delta^4)$$
$$\delta^4z_3^4(1-z_3)\arctan\frac{z_3^2-\delta^2}{\delta^2-4z_3+4}+(32768\,z_3^7-90112\,z_3^8+20480\,z_3^9+180224\,z_3^{10}-$$
$$251904\,z_3^{11}+140800\,z_3^{12}-35072\,z_3^{13}+3072\,z_3^{14}-258048\,z_3^5\,\delta^2+806912\,z_3^6\,\delta^2-845824\,z_3^7\,\delta^2+$$
$$87040\,z_3^8\,\delta^2+484864\,z_3^9\,\delta^2-382208\,z_3^{10}\,\delta^2+116096\,z_3^{11}\,\delta^2-12608\,z_3^{12}\,\delta^2-48\,z_3^{13}\,\delta^2+$$
$$49152\,z_3^3\,\delta^4-286720\,z_3^4\,\delta^4+642048\,z_3^5\,\delta^4-845824\,z_3^6\,\delta^4+871680\,z_3^7\,\delta^4-684544\,z_3^8\,\delta^4+$$
$$341824\,z_3^9\,\delta^4-90112\,z_3^{10}\,\delta^4+9712\,z_3^{11}\,\delta^4-288\,z_3^{12}\,\delta^4-20480\,z_3\,\delta^6+122880\,z_3^2\,\delta^6-$$
$$289280\,z_3^3\,\delta^6+453120\,z_3^4\,\delta^6-582144\,z_3^5\,\delta^6+489856\,z_3^6\,\delta^6-242112\,z_3^7\,\delta^6+64864\,z_3^8\,\delta^6-$$
$$6352\,z_3^9\,\delta^6-1336\,z_3^{10}\,\delta^6-138\,z_3^{11}\,\delta^6-12288\,\delta^8+36352\,z_3\,\delta^8-70400\,z_3^2\,\delta^8+136448\,z_3^3\,\delta^8-$$
$$160640\,z_3^4\,\delta^8+127616\,z_3^5\,\delta^8-62400\,z_3^6\,\delta^8+12272\,z_3^7\,\delta^8+1256\,z_3^8\,\delta^8-6\,z_3^9\,\delta^8+15\,z_3^{10}\,\delta^8-$$
$$768\,\delta^{10}-10752\,z_3\,\delta^{10}+29568\,z_3^2\,\delta^{10}-36480\,z_3^3\,\delta^{10}+18368\,z_3^4\,\delta^{10}-2144\,z_3^5\,\delta^{10}-472\,z_3^6\,\delta^{10}-$$
$$24\,z_3^7\,\delta^{10}+15\,z_3^8\,\delta^{10}-1152\,\delta^{12}+3072\,z_3\,\delta^{12}-2592\,z_3^2\,\delta^{12}+1344\,z_3^3\,\delta^{12}-696\,z_3^4\,\delta^{12}+$$
$$240\,z_3^5\,\delta^{12}-30\,z_3^6\,\delta^{12})\delta^2\sqrt{1-z_3}\ln\frac{z_3\sqrt{\delta^2-4z_3+4}+2\sqrt{(1-z_3)(z_3^2-\delta^2)}}{z_3\sqrt{\delta^2-4z_3+4}-2\sqrt{(1-z_3)(z_3^2-\delta^2)}}\Big). \qquad \text{(A2)}$$

[1] G.T. Bodwin, E. Braaten, and G.P. Lepage, Phys. Rev. **D51**, 1125 (1995); **D55**, 5853(E) (1997) (hep-ph/9407339).

[2] N. Brambillla *et al.* [quarkonium Working Group], hep-ph/0412158.

[3] K. Abe *et al.* [BELLE Collaboration], Phys. Rev. Lett. **89**, 142001 (2002) (hep-ex/0205104).

[4] P. Pakhlov, hep-ex/0412041.

[5] B. Aubert *er al.*[BABAR Collaboration], Phys. Rev. **D72**, 031101 (2005).

[6] E. Braaten and J. Lee, Phys. Rev. **D67**, 054007 (2003)(hep-ph/0211085); .

[7] Kui-Yong Liu, Zhi-Guo He, and Kuang-Ta Chao, Phys. Lett. **B557**, 45 (2003) (hep-ph/0211181) (see also, K.Y. Liu, Z.G. He, and K.T. Chao, hep-ph/0408141).

[8] K. Hagiwara, E. Kou, and C.-F.Qiao, Phys. Lett. **B570**, 39 (2003) (hep-ph/0305102).

[9] Y.J. Zhang, Y.J. Gao, and K.T. Chao, Phys. Rev. Lett.**96**, 092001 (2006) (hep-ph/0506076).

[10] D. Ebert and A.P. Martynenko, Phys. Rev. **D74**, 054008 (2006) (hep-ph/0605230).

[11] J.P. Ma and Z.G. Si, Phys.Rev. **D70** 074007(2004) (hep-ph/0405111); hep-ph/0608221.

[12] A.E. Bondar and V.L. Chernyak, Phys. Lett. **B612**, 215 (2005).(hep-ph/0412335)

[13] V.V. Braguta, A.K. Likhoded and A.V. Luchinsky, Phys. Rev. D72, 074019 (2005); hep-ph/0602047; hep-ph/0611021.

[14] V.V. Braguta, hep-ph/0701234.

[15] T. Huang and F. Zuo, hep-ph/0702147.

[16] G.T. Bodwin, D. Kang and J. Lee, Phys. Rev.**D74**, 014014 (2006) (hep-ph/0603186); hep-ph/0603185; G.T. Bodwin, D. Kang, D. Kim, J. Lee, and C. Yu, hep-ph/0611002.

[17] P. Cho and A.K. Leibovich, Phys. Rev. **D54** , 6690 (1996) (hep-ph/9606229).

[18] F. Yuan, C.F. Qiao, and K.T. Chao, Phys. Rev. **D56**, 321 (1997); *ibid*, 1663 (1997).

[19] S. Baek, P. Ko, J. Lee, and H.S. Song, J. Korean Phys. Soc. **33**, 97 (1998) (hep-ph/9804455).

[20] V.V. Kiselev, A.K. Likhoded, M.V. Shevlyagin, Phys. Lett. **B332**, 411 (1994)(hep-ph/9408407).

[21] Kui-Yong Liu, Zhi-Guo He, and Kuang-Ta Chao, Phys. Rev. **D69** (2004) 094027 (hep-ph/0301218).

[22] Kui-Yong Liu, Zhi-Guo He, and Kuang-Ta Chao, Phys. Rev. **D68**, 031501(R) (2003) (hep-ph/0305084).

[23] Yu-Jie Zhang and Kuang-Ta Chao, Phys. Rev. Lett. 98, 092003 (2007) (hep-ph/0611086).

[24] J.H. Kühn, J. Kaplan, and E.G. Safiani, Nucl. Phys. **B157**, 125 (1979); B. Guberina, J.H. Kühn, R.D. Peccei, and R. Rückl, Nucl. Phys. **B174**, 317 (1980); E.L. Berger and D.L. Jones, Phys. Rev. D23, 1521 (1981).

[25] G.T. Bodwin and A. Petrelli, Phys. Rev. **D66**, 094011 (2002) (hep-ph/0205210).

[26] Wai-Yee Keung and I.J. Muzinich, Phys. Rev. **D27**, 1518 (1983).

[27] E. Braaten and Y.Q. Chen, Phys. Rev. **D54**, 3216 (1996)(hep-pn/9604237).

[28] P. Ko, J. Lee, and H.S. Song, Phys. Rev **D54**, 4312 (1996) (hep-ph/9602223).

[29] W.-M.Yao *et al.* [Particle Data Group], J. Phys. **G33**, 1 (2006).

[30] M. Gremm, and A. Kapustin, Phys. Lett. **B407**, 323 (1997) (hep-ph/9701353).

[31] B.L. Ioffe and D.E. Kharzeev, Phys. Rev. **D69**, 014016 (2004) (hep-ph/0306062).

[32] A.V. Berezhnoy and A.K. Likhoded, Phys. Atom. Nucl. **67**, 757 (2004); Yad. Fiz **67** (2004) (hep-ph/0303145).

[33] A. B. Kaidalov, JETP Lett. **77**, 349 (2003).

[34] D. Kang, J. W. Lee, J. Lee, T. Kim and P. Ko, Phys. Rev. D **71**, 094019 (2005).